

## **General Disclaimer**

### **One or more of the Following Statements may affect this Document**

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

MLM-2906(OP)  
Conf-820325--4

**DISCLAIMER**  
This book was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

**MASTER**

MLM--2906 (OP)

**HYDROGEN DIFFUSION AND ELECTRONIC STRUCTURE**

DE82 008460

**IN CRYSTALLINE AND AMORPHOUS  $TiCu_y$**

R. C. Bowman, Jr.\*  
Division of Chemistry and  
Chemical Engineering  
California Institute of  
Technology  
Pasadena, California 91125

A. J. Maeland  
Materials Research Center  
Allied Chemical Corporation  
Morristown, New Jersey 07960

W.-K. Rhim  
Jet Propulsion Laboratory  
California Institute of  
Technology  
Pasadena, California 91103

J. F. Lynch\*\*  
Brookhaven National Laboratory  
Upton, New York 11973

**ABSTRACT**

Hydrogen diffusion behavior and electronic properties of crystalline  $TiCu_{0.94}$ ,  $Ti_2CuH_{1.90}$ , and  $Ti_2CuH_{2.63}$  and amorphous  $a-TiCuH_{1.4}$  are studied using proton relaxation times, proton Knight shifts, and magnetic susceptibilities. Crystal structure and hydrogen site occupancy have major roles in hydrogen mobility. The density of electron states at  $E_F$  is reduced in amorphous  $a-TiCuH_{1.4}$  compared to the crystalline hydrides.

The crystalline intermetallics  $TiCu$  and  $Ti_2Cu$  and the amorphous  $Ti_{1-y}Cu_y$  ( $0.3 \leq y \leq 0.7$ ) alloys directly react with gaseous hydrogen to form crystalline and amorphous ternary hydrides,<sup>1-4</sup> respectively, providing the temperature is maintained below 200°C. A recent nuclear magnetic resonance (NMR) study of the proton relaxation times<sup>5</sup> indicated a much higher hydrogen mobility in amorphous  $a-TiCuH_{1.3}$  compared

\*On leave from current address: Monsanto Research Corporation, Mound, Miamisburg, Ohio 45342.

\*\*Current address is Materials Research Center, Allied Chemical Corporation, Morristown, New Jersey 07960.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

MOU

to polycrystalline  $\text{TiCuH}_{0.94}$ . The increased disorder<sup>3</sup> of interstitial hydrogen occupancy in  $\alpha\text{-TiCuH}_x$  has been suggested<sup>5</sup> for the enhanced mobility in the amorphous phase. The present paper describes further NMR studies of diffusion in crystalline and amorphous  $\text{TiCuH}_x$  as well as crystalline  $\text{Ti}_2\text{CuH}_x$ . Furthermore, the electronic structure has been investigated using magnetic susceptibility, proton spin-lattice relaxation time ( $T_1$ ), and proton Knight shift ( $\sigma_K$ ) measurements.

Table 1 summarizes the structural properties of the  $\text{TiCuH}_x$  and  $\text{Ti}_2\text{CuH}_x$  samples as deduced from x-ray diffraction, neutron scattering, and proton lineshape parameters. The preparation procedures have been previously described.<sup>1-5</sup>

Table 1. Descriptions of  $\text{TiCuH}_x$  and  $\text{Ti}_2\text{CuH}_x$  and Hydrogen Diffusion Activation Energies ( $E_a$ )

| Sample                         | Metal Sublattice Structure | Probable Hydrogen Site Occupancies   | $E_a$ (eV)       | Temperature Range (K) |
|--------------------------------|----------------------------|--|------------------|-----------------------|
| $\text{TiCuH}_{0.94}$          | Tetragonal                 | 94% $\text{Ti}_4$ only   | $0.84 \pm 0.02$  | 465 - 560             |
| $\alpha\text{-TiCuH}_{1.4}$    | Amorphous                  | Mixed (mostly $\text{Ti}_4$ with some $\text{Ti}_4\text{-yCu}_y$ and octahedral) | $0.45 \pm 0.02$  | 357 - 410             |
|                                |                            |  | $0.185 \pm 0.01$ | 208 - 357             |
|                                |                            |  | $0.09 \pm 0.01$  | 150 - 207             |
| $\text{Ti}_2\text{CuH}_{1.9}$  | Orthorhombic (?)           | ~95% $\text{Ti}_4$ (some $\text{Ti}_2\text{Cu}_4$ likely)                        | $0.35 \pm 0.02$  | 290 - 519             |
| $\text{Ti}_2\text{CuH}_{2.63}$ | Orthorhombic (?)           | 100% $\text{Ti}_4$ and 63% $\text{Ti}_2\text{Cu}_4$                              | $0.29 \pm 0.02$  | 290 - 395             |

Hydrogen diffusion behavior has been evaluated using the temperature dependence of the proton rotating-frame spin-lattice relaxation times<sup>6</sup> ( $T_{1\rho}$ ) where the spin-locking field was about 7.3G and the proton resonance frequency was 34.5 MHz. The temperature dependences of the  $T_{1\rho}$  data for  $\text{Ti}_x\text{CuH}_x$  are shown in Fig. 1. Table 1 summarizes the diffusion activation energies ( $E_a$ ) that have been deduced from the  $T_{1\rho}$  data. Although a single  $E_a$  corresponding to Arrhenius behavior represents proton mobility in the crystalline  $\text{TiCuH}_{0.94}$  and  $\text{Ti}_2\text{CuH}_x$ , three  $E_a$  values are required for amorphous  $\alpha\text{-TiCuH}_{1.4}$ , which confirms the behavior previously seen<sup>5</sup> in  $\alpha\text{-TiCuH}_{1.3}$ . Furthermore,  $E_a$  is greatly reduced when protons occupy sites in addition to the tetrahedral  $\text{Ti}_4$  interstitials. This effect is seen in both crystalline  $\text{Ti}_2\text{CuH}_x$  and amorphous  $\alpha\text{-TiCuH}_{1.4}$ . From a consideration of the  $\text{TiCu}$  and  $\text{Ti}_2\text{Cu}$  crystal structures,<sup>1,4</sup> hydrogen diffusion in crystalline  $\text{TiCuH}$  can only occur by nearest neighbor jumps between the  $\text{Ti}_4$  sites,

while jumps through the intermediate  $\text{Ti}_2\text{Cu}_x$  sites become possible in  $\text{Ti}_2\text{CuH}_x$ . This probably accounts for the lower  $E_f$  values for  $\text{Ti}_2\text{CuH}_x$  and similar (or even easier) jump-paths are available in the more disordered amorphous phase.

The magnetic susceptibilities ( $\chi_m$ ) for  $\text{Ti}_y\text{CuH}_x$  were measured between 80 K and 300 K and are summarized in Fig. 2. Although the  $\chi_m$  values in Fig. 2 have been extrapolated to infinite magnetic field, the field-dependent ferromagnetic contribution was negligible except for a- $\text{TiCuH}_{1.4}$ , which appears to have some magnetic impurities as well as an opposite temperature dependence for  $\chi_m$ . There are several contributions to  $\chi_m$ , but only the paramagnetic term  $\chi_p$  is directly related to  $N(E_F)$ , the density of electron states at the Fermi level  $E_F$ . Hence, caution should be exercised in correlating  $\chi_m$  differences only to  $N(E_F)$  changes. In particular, the larger  $\chi_m$  for a- $\text{TiCuH}_{1.4}$  compared to  $\text{TiCuH}_{0.34}$  probably reflects either ferromagnetic or orbital contributions, and not a greater  $N(E_F)$  for the amorphous phase. However, the unusual<sup>7,8</sup>  $\chi_m$  increase with hydrogen content for  $\text{Ti}_2\text{CuH}_x$  is believed to actually correspond to  $N(E_F)$  becoming larger since the proton  $T_1$  and  $\sigma_K$  parameters also indicate  $N(E_F)$  increasing from  $\text{Ti}_2\text{CuH}_{1.9}$  to  $\text{Ti}_2\text{CuH}_{2.63}$ .




Fig. 2. Magnetic susceptibility values for  $\text{TiCuH}_x$  and  $\text{Ti}_2\text{CuH}_x$ .

The proton  $T_1$  and  $\sigma_K$  were measured at 56.4 MHz using methods previously described.<sup>9</sup> The  $\sigma_K$  values are referenced to tetramethylsilane. Table 2 summarizes the  $\sigma_K$  and  $(T_1 \cdot T)^{-1/2}$  parameters, which are directly proportional<sup>6,8,9</sup> to  $N(E_F)$ , at the upper and lower temperature limits of the present NMR measurements. The negative  $\sigma_K$  values in Table 2 indicate that core-polarization<sup>6</sup> with d-electrons dominates proton hyperfine interactions in  $Ti_xCuH_x$  where the population of d-states is much larger than s-states as has been previously found in other Ti-based hydrides.<sup>8,9</sup> Furthermore, the proton parameters suggest  $N(E_F)$  is significantly reduced in  $\alpha$ - $TiCuH_{1.4}$  compared to crystalline  $TiCuH_{0.94}$ , while  $N(E_F)$  increases with content in crystalline  $Ti_2CuH_x$ . However, a more detailed analysis based upon generalized Korringa relations<sup>6</sup> shows increased s-electron contact hyperfine interactions in the  $Ti_2CuH_x$  samples.

More extensive discussions of hydrogen diffusion and the electronic structures of  $Ti_xCuH_x$  will be published elsewhere.

Table 2. Proton Parameters  $(T_1 \cdot T)^{-1/2}$  and Knight Shifts  $\sigma_K$

| Sample                   | T<br>(K) | $(T_1 \cdot T)^{-1/2}$<br>(sK) <sup>-1/2</sup> | $\sigma_K$<br>(ppm) |
|--------------------------|----------|--|---------------------|
| $TiCuH_{0.94}$           | 300      | 0.163  | -120                |
|                          | 80       | 0.150  | -107                |
| $\alpha$ - $TiCuH_{1.4}$ | 210      | 0.113  | - 77                |
|                          | 80       | 0.108  | - 87                |
| $Ti_2CuH_{1.9}$          | 300      | 0.118  | - 67                |
|                          | 115      | 0.115  | - 69                |
| $Ti_2CuH_{2.63}$         | 300      | 0.140  | - 85                |
|                          | 80       | 0.145  | - 91                |

#### ACKNOWLEDGEMENTS

This work was partially supported by the Division of Chemical Sciences, Office of Basic Energy Sciences, U. S. Department of Energy, and the Caltech's President's Fund. Mound is operated by Monsanto Research Corporation for the U. S. Department of Energy under Contract No. DE-AC04-76-DP00053. Brookhaven National Laboratory is operated for the U. S. Department of Energy under Contract No. DE-AC-02-76-CH00016. Jet Propulsion Laboratory is operated for the National Aeronautics and Space Administration under Grant No. NAS7-100.

## REFERENCES

1. A. Santoro, A. Maeland, and J. J. Rush, Acta Cryst. B34: 3059 (1978).
2. A. J. Maeland, L. E. Tanner and G. G. Libowitz, J. Less-Common Met. 74: 279 (1980).
3. J. J. Rush, J. M. Rowe and A. J. Maeland, J. Phys. F: Metal Phys. 10: L283 (1980).
4. A. J. Maeland and G. G. Libowitz, J. Less-Common Met. 74: 295 (1980).
5. R. C. Bowman, Jr. and A. J. Maeland, Phys. Rev. B24: 2328 (1981).
6. R. M. Cotts, in "Hydrogen in Metals I: Basic Properties," G. Alefeld and V. Völkl, Ed., Springer-Verlag, Berlin, 1978, p. 227.
7. J. F. Lynch, R. Lindsay and R. O. Moyer, Jr., Solid State Commun. 41: 9 (1982).
8. J. F. Lynch, J. R. Johnson and R. C. Bowman, Jr., Paper at this meeting.
9. R. C. Bowman, Jr. and W.-K. Rhim, Phys. Rev. B24: 2232 (1981).

**- END -**

**DATE FILMED**

**4 / 16 / 82**




Fig. 1. Proton  $T_{1\rho}$  relaxation times with  $H_1 = 7.3$  G at  $\nu_H = 34.5$  MHz.